

HIGHLIGHT ARTICLE

Biomarkers in Neuroendocrine Tumors

Highlights from the "2013 ASCO Annual Meeting". Chicago, IL, USA; May 30 - June 4, 2013

Marvin Duque¹, Irvin M Modlin², Anumeha Gupta¹, Muhammad Wasif Saif¹

¹Division of Hematology Oncology, Tufts Cancer Center. Boston, MA, USA.

²Department of Surgery, Yale University School of Medicine. New Haven, CT, USA

Summary

Neuroendocrine tumors are a heterogeneous group of tumors with cells of neuroendocrine differentiation that arise from diverse anatomic sites with varying morphologic and clinical features. Since the natural history and prognosis varies widely between individual neuroendocrine tumor types, there is a critical need to identify accurate prognostic and predictive biomarkers and markers predictive of therapeutic efficacy. To date, plasma chromogranin-A levels have generally been accepted as the most useful biomarker, despite the fact that there are substantial concerns in sensitivity and discrepancies in measurement techniques. As a consequence, considerable attention has been focused upon the development of novel biomarkers that can be utilized with more clinical efficacy than chromogranin-A. In addition to amplifying the diagnostic/prognostic landscape, the need to calibrate the efficacy of biological targeted therapy has further accelerated the development of molecular biomarkers. At the 2013 American Society of Clinical Oncology (ASCO) Annual Meeting, Chou *et al.* (Abstract #e15151) presented data that chromogranin A levels can be monitored during treatment to predict clinical outcome. Modlin *et al.* (Abstract #4137), demonstrated a promising novel biomarker, serum multi-transcript molecular signature. Grande *et al.* (Abstract #4140), Heetfield *et al.* (Abstract #e15071) and Casanovas *et al.* (Abstract #4139) described sVEGFR2, p-mTOR and IGF1R as molecular markers with potential for use in targeted therapy trials. The authors review and summarize these abstracts in this article.

What Did We Know Before the 2013 American Society of Clinical Oncology (ASCO) Annual Meeting?

Neuroendocrine tumors (NETs) are a diverse family of tumors originating in neuroendocrine cells distributed ubiquitously throughout the body. The most common NETs are gastroenteropancreatic (GEP)-NETs and carcinoids. The incidence is 5.25 cases per 100,000 persons and the prevalence may exceed 100,000 in the United States [1, 2]. They

exhibit protean symptomatology and a spectrum of biological behavior [3]. The widely divergent biological behavior of individual NETs reflects their origin from a variety of neuroendocrine cell types (e.g., beta, G, EC and ECL cells) [4]. As a consequence, response to treatment and outcomes also vary considerably. For instance, stage IV gastrointestinal NETs exhibit a 5-year survival of 40.3% [5] compared to 15.5% for pancreatic NETs [1]. Current strategies in predicting prognosis include staging at surgery, histopathological assessment, Ki-67 index and grading [6]. Of particular relevance is the utility of biomarkers such as chromogranin A (CgA) and specific hormone products (insulin, gastrin, serotonin, etc.) of functional NETs, though the latter have become less used in recent times as CgA has been regarded as the most useful single measurement [3].

Chromogranin A levels

Chromogranin A levels, which represents a constitutive neuroendocrine secretory protein, is the most widely accepted biomarker. CgA was compared to a panel of biomarkers and proved to be the most accurate, with a specificity of 85.7% and sensitivity of 67.9% [7]. CgA is elevated in 60-

Key words Biological Markers; Chromogranin A; Eukaryotic Initiation Factor-4E; Genetic Markers; Molecular Targeted Therapy; Neuroendocrine Tumors; Receptor, IGF Type 1; TOR Serine-Threonine Kinases; Vascular Endothelial Growth Factor Receptor-2

Abbreviations CgA: chromogranin-A; GEP: gastroenteropancreatic; IGF1R: insulin-like growth factor 1 receptor; mTOR: mammalian target of rapamycin; NCI: National Cancer Institute; NET: neuroendocrine tumor; p-eIF4E: phosphorylated eukaryotic translation initiation factor 4E; p-mTOR: phosphorylated mTOR; sVEGFR2: soluble receptor for vascular endothelial growth factor 2

Correspondence Marvin Duque
Division of Hematology Oncology Tufts Cancer Center; 800 Washington St., #245; Boston, MA 02111, USA
Phone: +1-617-519-7445; Fax: +1-617.636.8538
E-mail: dr.marvin.duque@hotmail.com

80% of patients with neuroendocrine tumors. Elevated CgA levels correlate with disease burden and poor outcomes [8]. Serial CgA testing is being investigated to determine treatment response. In pancreatic NETs, early declines in CgA levels during treatment with chemotherapy and everolimus were associated with improved prognoses [9, 10]. However, the utility of serial CgA for monitoring treatment response still remains to be prospectively established [11]. The National Cancer Institute (NCI) recommends that serial measurements of CgA be incorporated into prospective clinical trials [12].

Novel Biomarkers

There is widespread recognition that CgA exhibits significant limitations as a single moiety, with concerns regarding both sensitivity and specificity. Since elevated CgA levels are found in cardiac, endocrine and inflammatory diseases, its specificity has been challenged, with reports of 10-35% specificity in GEP-NETs [13]. CgA is not expressed by 25% of NETs, resulting in a lower

sensitivity [14]. As such, CgA has not yet been accepted by the FDA [15] and is a Category 3 (with major disagreement) National Comprehensive Cancer Network (NCCN) recommendation [16]. Thus, it has become axiomatic that other strategies are required and consequently alternative novel biomarkers are being examined.

Molecular Biomarkers

The “success” of everolimus and sunitinib in the treatment of pancreatic NETs has emphasized the strategy of targeting angiogenesis and the PI3K/AKT/mTOR pathway. The major challenge in the targeted era is not only the varying responses to biological agents seen among diverse NETs but the need to develop techniques to predict efficacy [17]. Potential molecular biomarkers are therefore a source of substantial investigation. At this time, despite a paucity of rigorous evidence, the NCI recommends measuring methylguanine DNA methyltransferase (MGMT) expression to predict response in trials of alkylating agents and obtaining

Table 1. Abstracts presented at the 2013 ASCO Annual Meeting addressing neuroendocrine tumor biomarkers.

Biomarker	Study design	Results	Implications
CHROMOGRANIN A (CgA)			
CgA (Abstract #e15151) [18]	Retrospective n=54 with advanced GEP-NETs	Baseline CgA: Elevated in 85% Lower baseline CgA (<2 xULR): longer overall survival Serial CgA: Smaller increases (<15%); favorable response	Baseline and serial CgA may have utility to monitor outcomes during treatment
NOVEL BIOMARKERS			
Multi-transcript molecular signature (Abstract #4137) [19]	PCR score in training set (n=130) and validation set (n=182) of blood samples in GEP-NETs	PCR scores: Treatment naïve (vs. treated): higher (P<0.0001) Pre-treatment (vs. post-): reduced (P<0.004) Complete remission (vs. controls): not different Progressive disease (vs. stable): higher (P<0.0001) Identification metrics: Sensitivity 85-98%, specificity 93-97%, PPV 95-96%, NPV 87-98% Differentiation metrics: Sensitivity 91%, specificity 91%, PPV 86%, NPV 95%	Biomarker transcript panel very accurately identifies GEP-NETs, differentiates stable from progressive disease and exhibits potential to monitor treatment efficacy
MOLECULAR BIOMARKERS			
sVEGFR2 (Abstract #4140) [21]	Prospective Phase II n=44 with metastatic NETs treated with pazopanib	sVEGFR2: After treatment: decreased (P<0.001) Duration of treatment: decreased (P=0.0046) Decreased sVEGFR2 >20% (vs. <20%): longer mean progression free survival (12.6 vs. 9.1 months; P=0.067)	sVEGFR2 has potential to monitor efficacy of treatment specifically with pazopanib and multi-kinase inhibitors
p-mTOR IGF1R (Abstract #4139) [22]	Immunohistochemistry on tumor samples n=69 with NETs	p-mTOR: Expressed in 85% IGF1R: Expressed in 66% p-mTOR and IGF1R: Both expressed in 16%	IGF1R and p-mTOR may have some utility in the identification of appropriate patients for trials with mTOR inhibitors
p-mTOR p-eIF4E (Abstract #e15071) [23]	Retrospective n=26 tissue samples with poorly differentiated (G3) GEP-NETs	p-mTOR: Expressed in 64% High expression (vs. low): shorter mean overall survival (9 vs. 25 months; P=0.026) p-eIF4E: Expressed in 24% High expression (vs. low): shorter mean overall survival (9 vs. 25 months; P=0.015)	p-mTOR and p-eIF4E levels correlate with survival and may have some utility in the identification of appropriate patients for trials with mTOR inhibitors

GEP-NET: gastroenteropancreatic neuroendocrine tumor; HR: hazard ratio; IGF1R: insulin-like growth factor 1 receptors; n: number; NET: neuroendocrine tumors; NPV: negative predictive value; p-eIF4E: phosphorylated eukaryotic translation initiation factor 4E (eIF4E); p-mTOR: phosphorylated mammalian target of rapamycin; PPV: positive predictive value; sVEGFR2: soluble receptors for vascular endothelial growth factor 2; ULR: upper limit of reference

perfusion CT scans to measure decreases in tumor blood flow in trials of angiogenesis inhibitors [12]. The development of accurate and specific molecular biomarkers to tailor various molecular therapies to different NETs populations is a major unmet need.

What Did We Learn at the 2013 ASCO Annual Meeting?

Chromogranin A Levels

Chou *et al.* (Abstract #e15151) measured plasma chromogranin A level and abdominal CT scan at baseline and every 3 to 6 months after starting treatment in a retrospective study of 54 Asian patients with advanced gastroenteropancreatic NETs. Eighty-five percent had elevated baseline CgA level. Patients with lower baseline CgA levels (less than twice the upper limit of reference) had longer overall survival compared to those with higher levels (hazard ratio (HR)=0.39, 95% confidence interval (CI): 0.15-0.99), even after adjustment for other variables. Serial CgA levels were evaluated in 28 patients with a total of 81




Figure 1. Pipeline used to derive a set of 51 markers that identify gastroenteropancreatic neuroendocrine tumors. Two GEP-NEN gene expression datasets were analyzed (GEP-NEN-A and GEP-NEN-B). 1) GEP-NEN-A: small intestinal tissue (n=3; macroscopically normal mucosa collected at surgery), primary GEP-NENs (n=6), and metastatic GEP-NENs (n=3). 2) GEP-NEN-B: normal ileal mucosa (n=6), primary midgut neuroendocrine neoplasias (n=3), and liver metastases (n=3).

Step 1. Gene co-expression networks inferred from GEP-NEN-A and GEP-NEN-B datasets are intersected, producing the GEP-NEN network. **Step 2.** Co-expression networks from neoplastic and normal tissue microarray datasets are combined to produce the normal and neoplastic networks. **Step 3.** Links present in normal and neoplastic networks are subtracted from the GEP-NEN network. **Step 4:** Concordantly regulated genes in GEP-NEN-A and GEP-NEN-B networks are retained; other genes are eliminated from the GEP-NEN network, producing the Consensus GEP-NEN network. **Step 5.** Upregulated genes in both the GEP-NEN-A and GEP-NEN-B dataset are mapped to the Consensus GEP-NEN network. **Step 6.** Topological filtering, expression profiling, and literature-curation of putative tissue-based markers, yielding 21 putative genes further examined by RT-PCR. **Step 7.** Identification of mutually upregulated genes in GEP-NEN blood transcriptome and GEP-NEN-A and GEP-NEN-B datasets, yielding 32 putative genes further examined by RT-PCR. **Step 8.** Literature-curation and cancer mutation database search, yielding a panel of 22 putative marker genes for further RT-PCR. (Adopted with permission from Modlin *et al.* 2013 [20]). GEP-NEN: gastroenteropancreatic neuroendocrine neoplasia

measurements. A lower increase of CgA levels (less than 15%) predicted favorable antitumor response with a sensitivity of 86% and specificity of 91%. The authors concluded that these results demonstrated that measuring baseline and serial CgA levels may be used to predict clinical outcome and tumor response [18] (Table 1).

Novel Biomarkers

Modlin *et al.* (Abstract #4137) presented a novel candidate blood biomarker, a multi-transcript (n=51 genes) molecular signature measured by a serum RT-PCR and analyzed using a novel series of mathematical algorithms (Figure 1). They compared it to plasma CgA levels at baseline and following treatment. PCR scores were determined for GEP-NETs (n=63; treatment-naïve: n=28) in a training set and validated in an independent set. In the treatment-naïve GEP-NETs group, the PCR score was significantly elevated ($P < 0.0001$) compared to the treated group. In the 11 matched pre- and post-treatment samples, PCR levels were significantly reduced ($P < 0.004$) post-treatment. In the independent validation set, the PCR scores for complete remission were not different to controls (0.5 ± 0.25 vs. 0.4 ± 0.16), while PCR scores for stable disease (0.63 ± 0.12) were lower than for non-responders (5.85 ± 0.34 ; $P < 0.002$). PCR scores in progressive disease were significantly higher than for stable disease (85% vs. 10%, $P < 0.0001$). For identification, the multi-transcript gene signature demonstrated a high sensitivity (85-98%), specificity (93-97%), positive predictive value (PPV) (95-96%) and negative predictive value (NPV) (87-98%). Likewise, in detecting differentiation between stable and progressive disease, the biomarker also exhibited high sensitivity (91%), specificity (91%), PPV (86%) and NPV (95%), significantly outperforming plasma CgA testing ($P < 0.005$). This study demonstrated that the MTMS is a promising novel multifactorial neuroendocrine specific biomarker to detect NETs with the potential for measuring response to treatment [19, 20] (Table 1).

Molecular Biomarkers

Grande *et al.* (Abstract #4140) assessed soluble receptor for vascular endothelial growth factor 2 (sVEGFR2) in predicting clinical outcome in patients with metastatic NETs treated with pazopanib, a multi-tyrosine kinase/VEGF inhibitor. Biomarkers were assessed at baseline and after 12 weeks of treatment. sVEGFR2 levels were found to be decreased at 12 weeks (median decrease: 20%, $P < 0.0001$), as well as with a longer duration of treatment ($P = 0.0046$). Patients with a greater decrease in sVEGFR2 (more than 20% decrease vs. less than 20%) trended towards an improved mean progression free survival (12.6 vs. 9.1 months) ($P = 0.067$) [21] (Table 1).

Casanovas *et al.* (Abstract #4139) described the activation status of insulin-like growth factor 1 receptors (IGF1R)-mTOR pathway by immunohistochemistry in 69 tumor samples of NETs. IGF1R was expressed in 66% and phosphorylated mTOR (p-mTOR) was expressed in 20% (lower than expected). Expression of both molecular markers, indicating consistent activation of the IGF1R-mTOR pathway, was only found in 16%. A subgroup of ileal NETs showed consistent activation of both IGF1R and p-mTOR [22] (Table 1).

Heetfeld *et al.* (Abstract #e15071) analyzed 26 cases of poorly differentiated GEP-NETs for molecular biomarkers p-mTOR and phosphorylated eukaryotic translation initiation factor 4E (p-eIF4E). p-mTOR was expressed in 64% and its downstream effector p-eIF4E was expressed in 24%. High expression levels of p-mTOR (mean overall survival: 9 vs. 25 months; P=0.026) and expression of p-eIF4E (mean overall survival: vs. 25 months; P=0.015) was significantly associated with shorter survival. Both biomarkers were also independent prognostic factors for survival when considering other factors [23] (Table 1).

Discussion

In neuroendocrine tumor management, biomarkers are of critical value since the use of imaging in a disease that is often indolent has significant limitations. In addition they have the potential to detect disease early, risk stratify patients and monitor response to treatment. Chromogranin A, a neuroendocrine secretory protein, has obvious biological limitations in predicting cell proliferation and therapeutic efficacy. Although it is a widely used biomarker, the limitations of a mono-marker are apparent and considerable further validation in measuring therapeutic response is required. In particular the dependence upon a single agent (CgA or pancreastatin) or minor combinations thereof represents a vulnerability when seeking to predict biological behavior that represents a summation of a multiplicity of gene regulated events that reflect cell proliferation, metastasis or drug sensitivity. As a consequence of these limitations, new biomarkers are being developed. The movement towards biological targeted therapy has emphasized the importance of identifying biomarkers that will enable clinicians to identify patients who will most likely respond to specific molecular agents or be able to define early evidence of therapeutic efficacy.

The 2013 ASCO Annual Meeting presented important advances in neuroendocrine biomarkers. Chou *et al.* (Abstract #e15151) showed that lower levels of CgA predicted significantly improved survival and a favorable response to treatment, providing evidence that baseline and serial CgA

levels may be used to monitor clinical outcome [18]. Modlin *et al.* (Abstract #4137) demonstrated that a multi-transcript molecular signature for PCR blood analysis outperformed CgA levels in identifying GEP-NETs and in detecting differences in response [19]. Grande *et al.* (Abstract #4140) noted that in patients treated with pazopanib, sVEGFR2 levels were decreased after treatment and a quantitatively larger decrease was associated with improved progression free survival [21]. sVEGFR2 may have potential as a molecular marker in VEGF and multi-kinase inhibitor therapy. Casanovas *et al.* (Abstract #4139) found that while a majority of NETs express IGF1R, only a small subset demonstrate activation of the IGF1R-mTOR pathway [22]. p-mTOR and IGF1R expression could potentially help stratify patients in future clinical trials. Heetfeld *et al.* (Abstract #e15071) showed that in poorly differentiated NETs, high expression of phosphorylated p-mTOR and its downstream effector, p-eIF4E was associated with a significantly shorter survival [23]. The identification of these p-mTOR and p-eIF4E raise the possibility, albeit limited, that in individuals with aggressive poorly differentiated NETs, mTOR inhibitors may be reasonable to evaluate in trials. However, given that mTOR inhibitors may be of limited value in indolent disease [24], this proposal may seem quixotic to some. Further studies are needed to validate these markers and define their clinical significance.

Conflict of interest The authors have no potential conflict of interest

References

1. Yao JC, Hassan M, Phan A, et al. One hundred years after "carcinoid": epidemiology of and prognostic factors for neuroendocrine tumors in 35,825 cases in the United States. *J Clin Oncol* 2008;26:3063-3072.
2. Tsikitis VL, Wetheim BC, Guerrero MA. Trends of incidence and survival of gastrointestinal neuroendocrine tumors in the United States: a SEER analysis. *J Cancer* 2012;3:292-302.
3. Modlin IM, Oberg K, Chung DC et al. Gastroenteropancreatic neuroendocrine tumors. *Lancet Oncol* 2008; 9:61-72
4. Lawrence B, Gustafsson BI, Chan A, et al. The epidemiology of gastroenteropancreatic neuroendocrine tumors. *Endocrinol Metab Clin North Am.* 2011 40(1):1-18.
5. Edge SR, Byrd DR, Compton CC et al. Neuroendocrine tumors. *AJCC Staging Handbook*, 7th edition. 2010.: 227-245.
6. Klimstra DS, Modlin IM, Coppola D, et al. The pathologic classification of neuroendocrine tumors: a review of nomenclature, grading, and staging systems. *Pancreas.* 2010;39(6):707-12.
7. Bajetta E, Ferrari L, Martinetti A, et al. Chromogranin A, Neuron Specific Enolase, Carcinoembryonic Antigen and Hydroxyindole Acetic Acid Evaluation in Patients with Neuroendocrine Tumors. *Cancer* 1999;86:858-865.
8. Moran CA, Suster S, Coppola D, et al: Neuroendocrine carcinomas of the lung: A critical analysis. *Am J Clin Pathol* 131:206-221, 2009

9. Yao JC, Lombard-Bohas C, Baudin E, et al: Daily oral everolimus activity in patients with metastatic pancreatic neuroendocrine tumors after failure of cytotoxic chemotherapy: A phase II trial. *J Clin Oncol* 28:69-76, 2010.
 10. Kouvaraki MA, Ajani JA, Hoff P, et al. Fluorouracil, doxorubicin, and streptozocin in the treatment of patients with advanced and metastatic pancreatic neuroendocrine carcinomas. *J Clin Oncol* 2004;22:4762-4771.
 11. Belli SH, Oneto A, Aranda C, et al. Chromogranin A as a biochemical marker for the management of neuroendocrine tumors: a multicenter study developed in Argentina. *Acta Gastroenterol Latinoam*. 2009 Sep;39(3):184-9.
 12. Kulke M, Sui L, Tepper J et al. Future Directions in the Treatment of Neuroendocrine Tumors: Consensus Report of the National Cancer Institute Tumor Clinical Trials Planning Meeting. *J Clin Oncol* 2011; 29:934-943.
 13. Modlin IM, Gustafsson BI, Moss SF, et. al. Chromogranin A--biological function and clinical utility in neuro endocrine tumor disease. *Ann Surg Oncol*. 2010;17(9):2427
 14. Zatelli MC, Torta M, Leon A et al. Chromogranin A as a marker of neuroendocrine neoplasia: an Italian Multicenter Study. *Endoc Relat Cancer* 2007; 14:473-482.
 15. AACC (2010) Chromogranin A. Lab Tests Online: American Association for Clinical Chemistry. Pp. Details Regarding Chromogranin A as a test.
 16. NCCN Clinical Practice Guidelines in Oncology/ Neuroendocrine Tumors. Version 2.2013. http://www.nccn.org/professionals/physician_gls/neuroendocrine.pdf. Accessed June 6, 2013.
 17. Leung R, Lang B, Wong H et al. Advances in the systemic treatment of neuroendocrine tumors in the era of molecular therapy. *Anticancer Agents Med Chem* 2013;13(3):382-388.
 18. Chou WC and Chang L. Baseline and dynamic plasma chromogranin A (CgA) level to predict clinical outcome and tumor response in Asian patients with gastroenteropancreatic neuroendocrine tumor (GEP-NET). *J Clin Oncol* 31, 2013 (suppl; abstr #e15151).
 19. Modlin IM, Drozdov I and Kidd, MS. A multi-transcript blood neuroendocrine tumor molecular signature to identify treatment efficacy and disease progress. *J Clin Oncol* 31, 2013 (suppl; abstr #4137).
 20. Modlin IM, Drozdov I, Kidd M. The identification of gut neuroendocrine tumor disease by multiple synchronous transcript analysis in blood. *PLoS One*. 2013 15;8(5):e63364.
 21. Grande E, Casanovas O, Earl J, et al. sVEGFR2 and circulating tumor cells to predict for the efficacy of pazopanib in neuroendocrine tumors (NETs): PAZONET subgroup analysis. *J Clin Oncol* 31, 2013 (suppl; #abstr 4140).
 22. Casanovas O, Teule A, Villabona C, et al. Study on activation of the IGF-1R mTOR pathway in neuroendocrine tumours (NETs). *J Clin Oncol* 31, 2013 (suppl; abstr #4139).
 23. Heetfeld M, Kasajima A, Koch A, et. al. Poorly differentiated neuroendocrine carcinoma (NEC G3): Prognostic factors and potential novel targets. *J Clin Oncol* 31, 2013 (suppl; abstr #e15071).
 24. Pavel ME, Hainsworth JD, Baudin E, et. al. RADIANT-2 Study Group. Everolimus plus octreotide long-acting repeatable for the treatment of advanced neuroendocrine tumours associated with carcinoid syndrome (RADIANT-2): a randomised, placebo-controlled, phase 3 study. *Lancet*. 2011 10;378(9808):2005-12.
-